

On finite-dimensional absolute-valued algebras satisfying $(x^p, x^q, x^r) = 0$

A. Chandid, M. I. Ramírez and A. Rochdi

February 11, 2010

Abstract

By means of principal isotopes $\mathcal{H}(a, b)$ of algebra $\mathcal{H}$ [Ra 99] we give an exhaustive description of all 4-dimensional absolute valued algebras satisfying $(x^p, x^q, x^r) = 0$ for fixed integers $p, q, r \in \{1, 2\}$. For such an algebras the number $N(p, q, r)$ of isomorphisms classes is either finite included between two and three, or is infinite. Concretely

1. $N(1, 1, 1) = N(1, 1, 2) = N(1, 2, 1) = N(2, 1, 1) = 2$,
2. $N(1, 2, 2) = N(2, 2, 1) = 3$,
3. $N(2, 1, 2) = N(2, 2, 2) = \infty$.

Besides, each one of above algebras contains 2-dimensional subalgebras. However, the problem in dimension 8 is far from being completely solved. In fact, there are 8-dimensional absolute-valued algebras, containing no 4-dimensional subalgebras, satisfying $(x^2, x, x^2) = (x^2, x^2, x^2) = 0$.

Keywords. Absolute valued algebra, central (flexible) idempotent, principal isotopes of $\mathcal{H}$.

1 Introduction

Absolute-valued algebras, introduced in 1918 [Ost 18], constitute nowadays a distinguished and attractive class of non-associative algebras, by its widely observed variety, and we find in [Rod 04] a perfect compilation of the results around the theory, previous in 2004. During the last decade several works on finite-dimensional

absolute-valued algebras appeared [Ra 99], [Roc 03], [Rod 04], [CM 05], [CR 08], [RR 09], [CKMMRR]. A fundamental result on this subject, given by Albert [A 47], asserts that the dimension of every finite-dimensional absolute-valued algebra A is 1, 2, 4 or 8, and A is isotopic to one of classical (unital) absolute-valued algebras $\mathbb{R}$, $\mathbb{C}$, $\mathbb{H}$ (real quaternion algebra) or $\mathbb{O}$ (real octonion algebra). It follows easily that $\mathbb{R}, \mathbb{C}, {}^*\mathbb{C}, \mathbb{C}^*, \mathbb{C}^*$ are the unique absolute-valued real algebras of dimension ≤ 2 . Using the so-called principal isotopes of $\mathbb{H}$ formed by four families depending on a pair of norm-one elements of $\mathbb{H}$, Ramírez [Ra 99] classified all 4-dimensional absolute-valued real algebras and solved the isomorphism problem. Calderón and Martín [CM 05] gave a refinement of above classification. Recently a description of all 8-dimensional absolute-valued algebras was given [CKMMRR]. On the other hand, classical results specify those absolute-valued algebras which satisfy to a familiar identity as associativity [Ost 18], [M 38] or commutativity [UW 60] and where extended to power-associativity [EM 80] and flexibility [EM 81] respectively. Concretely

1. Every absolute-valued associative real algebra is isometrically isomorphic to either $\mathbb{R}, \mathbb{C}$ or $\mathbb{H}$.
2. $\mathbb{R}, \mathbb{C}$, and $\mathbb{C}^*$ are the unique absolute-valued commutative real algebras.
3. $\mathbb{R}, \mathbb{C}, \mathbb{H}$, and $\mathbb{O}$ are the unique absolute-valued power-associative real algebras.
4. Every absolute-valued flexible real algebra is finite-dimensional.

By considering third power-associative algebras, that is algebras satisfying $(x, x, x) = 0$ where $(., ., .)$ means associator, El-Mallah [Elm 87] and, independently, Elduque-Pérez [EP 94] gave a finite-dimensional extension of above results by showing that every finite-dimensional absolute-valued algebra satisfying $(x, x, x) = 0$ is flexible and equal to either $\mathbb{R}, \mathbb{C}, \mathbb{C}^*, \mathbb{H}, \mathbb{H}^*, \mathbb{O}, \mathbb{O}^*$, or the algebra $\mathbb{P}$ of pseudo-octonions. In a general way absolute-valued algebras satisfying to an identity of the form $(x^p, x^q, x^r) = 0$ for fixed integers $p, q, r \in \{1, 2\}$, including $(x, x, x) = 0$, where extensively studied [Elm 82, 83, 87, 88, 97, 01], [EP 94], [EE 04], [Cu 06], [CR 08], [RR 09], [MRR]. For such an algebra A , the existence of either a non-zero central idempotent [Elm 83, 01], [EE 04], [Cu 06], [CR 08], [RR 09], or a left-unit [CR 08] carry away finite dimensionality. Moreover, a complete description of these algebras was given in both cases ([Elm 88], [Cu 06], [CR 08], [RR 09]) and ([CR 08], [MRR]), respectively. However there are infinite-dimensional absolute-valued algebras satisfying to the identity $(x^2, y, x^2) = 0$, in two variables [CR 08]. In the other hand it

seems that, with the exception of identity $(x, x, x) = 0$, no previous study took place for an identity of the form $(x^p, x^q, x^r) = 0$ over an arbitrary absolute-valued algebra of finite dimension ≥ 4 . The aim of this paper is to make some contributions to this subject. Here we give a description of all four-dimensional absolute-valued algebras satisfying an identity of the form $(x^p, x^q, x^r) = 0$ for fixed integers $p, q, r \in \{1, 2\}$ (Theorem 3.24). We show the following main result:

Theorem. *Let A be a four-dimensional absolute valued algebra. Then A satisfies to an identity of the form $(x^p, x^q, x^r) = 0$, for fixed integers $p, q, r \in \{1, 2\}$, if and only if A is isomorphic to one of principal isotopes of $\mathbb{H}$ described by the following table:*

A satisfies to	The list of isomorphism classes
$(x, x, x) = 0$ or $(x, x^2, x) = 0$	$\mathbb{H}, {}^*\mathbb{H}$
$(x, x, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*$
$(x, x^2, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*, \mathbb{H}^*(1, i)$
$(x^2, x, x) = 0$	$\mathbb{H}, {}^*\mathbb{H}$
$(x^2, x^2, x) = 0$	$\mathbb{H}, {}^*\mathbb{H}, {}^*\mathbb{H}(i, 1)$
$(x^2, x, x^2) = 0$	$\mathbb{H}, {}^*\mathbb{H}$ ${}^*\mathbb{H}(1, e^{i\alpha}), \mathbb{H}^*(e^{i\alpha}, 1)$ with $\alpha \in [0, \pi[$
$(x^2, x^2, x^2) = 0$	$\mathbb{H}, {}^*\mathbb{H}$ ${}^*\mathbb{H}(i, 1), {}^*\mathbb{H}(i, i), \mathbb{H}^*(1, i), \mathbb{H}^*(i, i)$ ${}^*\mathbb{H}(1, e^{i\alpha}), \mathbb{H}^*(e^{i\alpha}, 1)$ with $\alpha \in [0, \pi[$

We can easily see that each of above algebras contains 2-dimensional subalgebras. However, the problem in dimension 8 is very hard. There are 8-dimensional absolute-valued algebra satisfying $(x^2, y, x^2) = 0$ containing no 4-dimensional subalgebras. It is well known that every third power-associative algebra, over a field of characteristic $\neq 2$ having at least three elements, satisfies to $(x, x^2, x) = 0$. In the other hand we see that, among those 4-dimensional absolute-valued algebras, the identity $(x, x^2, x) = 0$ carry away third power-associativity. However the question to know if an 8-dimensional absolute-valued algebra which satisfies both identities $(x, x^2, x) = (x^2, y, x^2) = 0$ is third power-associative, seems still to be an open problem.

2 Notations and preliminary results

Definitions and notations 2.1 *An algebra over an arbitrary field K is a vector space over K endowed with a bilinear mapping $(x, y) \mapsto xy$ from $A \times A$ to A called the product of the algebra A . For x, y, z in A , we denote by (x, y, z) the associator $(xy)z - x(yz)$ of x, y, z and by $[x, y]$ the commutator $xy - yx$ of x, y . We denote by L_x, R_x the linear operators of A defined by $y \mapsto xy, y \mapsto yx$ respectively. An element $e \in A$ is said to be **central** if it satisfies $[e, x] = 0$ for all $x \in A$, or, in other word $L_e = R_e$. The element $e \in A$ is said to be **flexible** if it satisfies $(e, x, e) = 0$ for all $x \in A$, or, in other words $L_e \circ R_e = R_e \circ L_e$.*

1. The algebra A is said to be

- (a) **Third power-associative** if it satisfies the identity $(x, x, x) = 0$.
- (b) **Alternative** if it satisfies both identities $(x, x, y) = (y, x, x) = 0$. A well known Artin's theorem asserts that an algebra is alternative if and only if the subalgebra generated by two arbitrary elements is associative ([Sc 66] Theorem 3.1 p. 29).
- (c) **Flexible** it satisfies the identity $(x, y, x) = 0$.
- (d) **Quadratic** if it contains an unit element e and e, x, x^2 are linearly dependent for all $x \in A$. The elements in the set

$$Im(A) = \{x \in A : x^2 \in Ke \text{ and } x \notin Ke - \{0\}\}$$

are called the purely imaginary elements in A . If, moreover, the characteristic of K is not two, then $Im(A)$ is a subspace of A supplementary to Ke , i.e., $A = Ke \oplus Im(A)$ ([HKR 91] p. 227-228).

2. A linear operation $*$: $x \mapsto x^*$ defined on algebra A is said to be an involution if it satisfies, for all $x, y \in A$, the following conditions

- (a) $(x^*)^* = x$,
- (b) $xx^* = x^*x$,
- (c) $(xy)^* = y^*x^*.$ $\square$

Proposition 2.2 *Let A be an algebra over a field K of characteristic $\neq 2$ containing at least three elements. If algebra A is third power-associative then A satisfies to the identity $(x, x^2, x) = 0$.*

Proof. A simple linearization of $(x, x, x) = 0$, taking into account that K contains two distinct non-zero elements, gives $[x^2, y] + [xy + yx, x] = 0$. The result follows by putting $y = x^2$ in above identity, taking into account the characteristic of K . $\square$

Definitions 2.3 *An algebra A , over real numbers $\mathbb{R}$ or complex numbers $\mathbb{C}$, is said to be absolute valued if the space A is endowed with a norm $\|\cdot\|$ satisfying $\|xy\| = \|x\| \|y\|$ for all $x, y \in A$. $\square$*

Remarks 2.4 *The algebras $\mathbb{C}$, $\mathbb{H}$ (quaternions) and $\mathbb{O}$ (octonions) can be built from $\mathbb{R}$ by iterating the Cayley-Dickson doubling process ([HKR 91] p. 256-257). If A stands for either $\mathbb{R}$, $\mathbb{C}$, $\mathbb{H}$ or $\mathbb{O}$, with unit element 1 and standard involution (SI) $\sigma_A : x \mapsto \bar{x}$, then*

1. *A is a quadratic and alternative algebra, and the expression $\mathbb{R} \oplus \text{Im}(A)$ is nothing other than the decomposition of A into a direct sum of the eigenspaces $E_1(\sigma_A)$, $E_{-1}(\sigma_A)$ associated, respectively, to eigenvalues 1 and -1 . Thus every element $x \in A$ can be written as a sum of its real part $\frac{1}{2}(x + \bar{x}) := \text{Re}(x) \in \mathbb{R}$ and purely imaginary part $\frac{1}{2}(x - \bar{x}) := \text{Im}(x) \in \text{Im}(A)$.*
2. *The usual inner product*

$$(\cdot|\cdot) : A \times A \rightarrow \mathbb{R}^+ \quad (x, y) \mapsto (x|y) = \text{Re}(x\bar{y})$$

([HKR 91] p. 208, 253) satisfies to $(xy|xy) = (x|x)(y|y)$ for all $x, y \in A$ and the (euclidian) norm $\|\cdot\|$ of A , given by $\|x\| = \sqrt{x\bar{x}} = \sqrt{\bar{x}x}$ for all $x \in A$, is an absolute-value. Thus $(A, \|\cdot\|)$ is a pre-Hilbert real absolute-valued algebra and the SI σ_A of $A = \mathbb{R} \oplus \text{Im}(A)$ is a linear isometry. Besides, the following equalities holds for all $x, y \in A$:

- (a) $(x|y) = (\bar{x}|\bar{y})$. *This follows from equality $(\bar{x}|\bar{x}) = (x|x)$ by linearization.*
- (b) $(\cdot|\cdot)$ is a trace form on A ([BBO 82] Lemma 1.2):

$$(xy|z) = (x|yz) \tag{2.1}$$

- (c) *The triple product identity ([HKR 91] p. 209, 254):*

$$yxy = 2(\bar{x}|y)y - \|y\|^2\bar{x}. \tag{2.2}$$

According to [A 47], two absolute-valued algebras A and B , are said to be isotopic if there exist linear isometries Φ_1, Φ_2, Φ_3 from A onto B satisfying $\Phi_1(xy) = \Phi_2(x)\Phi_3(y)$ for all $x, y \in A$. One of fundamental results about finite-dimensional absolute-valued algebras, proved by Albert ([A 47] Theorem 2), is the following:

Theorem 2.5 *Let A be a finite-dimensional absolute-valued real algebra. Then A is isotopic to either $\mathbb{R}, \mathbb{C}, \mathbb{H}$ or $\mathcal{O}$. Therefore A has dimension 1, 2, 4 or 8 and the absolute-value of A comes from an inner product.* $\square$

Examples 2.6 .

1. **(Standard isotopes of $\mathbb{A}$).** For $\mathbb{A}$ equal to either $\mathbb{C}, \mathbb{H}$ or $\mathcal{O}$ with SI $x \mapsto \bar{x}$, let us denote by ${}^*\mathbb{A}, \mathbb{A}^*$ and $\overset{*}{\mathbb{A}}$ the absolute-valued real algebras obtained by endowing the normed space $\mathbb{A}$ with the products $x \odot y = \bar{x}y$, $x \odot y = x\bar{y}$ and $x \odot y = \bar{x}\bar{y}$, respectively. Algebras $\mathbb{A}, {}^*\mathbb{A}, \mathbb{A}^*, \overset{*}{\mathbb{A}}$ are called standard isotopes of $\mathbb{A}$. It follows easily from Theorem 2.5 that $\mathbb{C}, {}^*\mathbb{C}, \mathbb{C}^*$ and $\overset{*}{\mathbb{C}}$ are the unique 2-dimensional absolute-valued real algebras (see [Rod 94], [EP 99]).
2. **(Principal isotopes of $\mathbb{H}$).** These are the absolute-valued algebras $\mathbb{H}_1(a, b), \mathbb{H}_2(a, b), \mathbb{H}_3(a, b)$ and $\mathbb{H}_4(a, b)$ obtained from fixed norm-one elements a, b in $\mathbb{H}$ by endowing the normed space $\mathbb{H}$ with the products

$$\begin{aligned} x \overset{a,b}{\odot}_1 x &= axyb, \\ x \overset{a,b}{\odot}_2 x &= \bar{x}ayb, \\ x \overset{a,b}{\odot}_3 y &= axb\bar{y}, \\ x \overset{a,b}{\odot}_4 y &= a\bar{x}\bar{y}b. \end{aligned}$$

These four types of absolute-valued algebras are called principal isotopes of $\mathbb{H}$ denoted $\mathbb{H}(a, b), {}^*\mathbb{H}(a, b), \mathbb{H}^*(a, b)$ and $\overset{*}{\mathbb{H}}(a, b)$, respectively [Ra 99]. $\square$

M. I. Ramírez [Ra 99] gave a classification of all 4-dimensional absolute-valued algebras by using Theorem 2.5 and the description of all linear isometries on $\mathbb{H}$ ([HKR 91] p. 215):

Theorem 2.7 *Every four-dimensional absolute valued real algebra is isomorphic to a principal isotope of $\mathbb{H}$. Moreover two principal isotopes $\mathbb{H}_n(a, b)$ and $\mathbb{H}_m(a', b')$ of $\mathbb{H}$ are isomorphic if and only if $n = m$ and the equalities $a'p = \varepsilon pa$ and $b'p = \delta pb$ hold for some norm-one element $p \in \mathbb{H}$ and some $\varepsilon, \delta \in \{1, -1\}$.* $\square$

Remarks 2.8 Let n be in $\{1, 2, 3, 4\}$.

1. Algebras $\mathbb{H}_n(-a, -b)$, $\mathbb{H}_n(-a, b)$, $\mathbb{H}_n(a, -b)$, $\mathbb{H}_n(a, b)$ are isomorphic.
2. A refinement of Theorem 2.7 can be found in [CM 05].
3. We check easily that, among those principal isotopes of $\mathbb{H}$, algebras $\mathbb{H}(a, \bar{a})$, ${}^*\mathbb{H}(a, \bar{a})$ are the unique ones which contain a non-zero central idempotent e , inevitably equal to 1. $\square$

Ramírez [Ra 99] gave, among others, a precise description of all 4-dimensional absolute-valued algebras which contain 2-dimensional sub-algebras:

Theorem 2.9 A 4-dimensional absolute-valued real algebra contains subalgebras of dimension 2 if and only if it is isomorphic to a principal isotope of $\mathbb{H}$ whose parameters a, b are precisely subjected to the conditions given by the following table

$\mathbb{H}(a, b)$	${}^*\mathbb{H}(a, b)$	$\mathbb{H}^*(a, b)$	${}^*\mathbb{H}(a, b)$	contains
$ab = ba$	$a, ba \in \text{Im}(\mathbb{H})$	$b, ba \in \text{Im}(\mathbb{H})$	$a, b \in \text{Im}(\mathbb{H})$	$\mathbb{C}$
$a, ab \in \text{Im}(\mathbb{H})$	$ab = ba$	$a, ba \in \text{Im}(\mathbb{H})$	$a, ba \in \text{Im}(\mathbb{H})$	${}^*\mathbb{C}$
$b, ba \in \text{Im}(\mathbb{H})$	$b, ba \in \text{Im}(\mathbb{H})$	$ab = ba$	$b, ba \in \text{Im}(\mathbb{H})$	$\mathbb{C}^*$
$a, b \in \text{Im}(\mathbb{H})$	$a, b \in \text{Im}(\mathbb{H})$	$a, b \in \text{Im}(\mathbb{H})$	$ab = ba$	$\mathbb{C}^*$

We shall need the following useful result:

Lemma 2.10 For a, b in $\mathbb{H}$, with the same norm, the the following are equivalents:

1. a and b are conjugated.
2. $\text{Re}(a) = \text{Re}(b)$.

Proof. The implication $\mathbf{1} \Rightarrow \mathbf{2}$ is clear and it remains only to show $\mathbf{2} \Rightarrow \mathbf{1}$ with $a, b \in \text{Im}(\mathbb{H})$. Indeed, we have: $b(a + b) = ba + b^2 = ba + a^2 = (a + b)a$.

1. If $b \neq -a$, then $b = va\bar{v}$ where $v = \|a + b\|^{-1}(a + b)$.
2. If $b = -a \neq 0$, then for all norm-one $u \in \text{Im}(\mathbb{H})$, orthogonal to a , we have:
 $b = u^2a = -uau = ua\bar{u}$. $\square$

3 Classification in dimension four

3.1 Principal isotopes of $\mathcal{H}$ satisfying to $(x^p, x^q, x^r) = 0$

Now we will study those finite-dimensional absolute-valued algebras which satisfy an identity of the form $(x^p, x^q, x^r) = 0$ for fixed integers $p, q, r \in \{1, 2\}$. The situation in dimension one, corresponding to the associative algebra $\mathbb{R}$, where all the identities $(x^p, x^q, x^r) = 0$ are satisfied, is trivial. It is not so much in dimension two:

Proposition 3.1 *Among all 2-dimensional absolute-valued algebras*

1. $\mathcal{C}$ is the unique which satisfies all the identities $(x^p, x^q, x^r) = 0$.
2. The other algebras, ${}^*\mathcal{C}, \mathcal{C}^*, \mathcal{C}^*$, satisfy to $(x^2, x, x^2) = (x^2, x^2, x^2) = 0$. Among these three last ones
 - (a) ${}^*\mathcal{C}$ is the unique which satisfy to $(x^2, x, x) = 0$ or else, to $(x^2, x^2, x) = 0$.
 - (b) $\mathcal{C}^*$ is the unique which satisfy to $(x, x, x^2) = 0$ or else, to $(x, x^2, x^2) = 0$.
 - (c) $\mathcal{C}^*$ is the unique which satisfy to $(x, x, x) = 0$ or else, to $(x, x^2, x) = 0$. $\square$

We will show that the general situation in dimension 4 is very diversified and principal isotopes of $\mathcal{H}$ will be used for our study. First of all the following remark, for standard isotopes of $\mathcal{H}$, will serve us as guide:

Remark 3.2 *Let us note that the associative algebra $\mathcal{H}$ satisfy to all the identities of the form $(x^p, x^q, x^r) = 0$ and that algebras ${}^*\mathcal{H}, \mathcal{H}^*, \mathcal{H}^*$ satisfy to both identities $(x^2, x, x^2) = (x^2, x^2, x^2) = 0$. That, among these last three algebras*

1. ${}^*\mathcal{H}$ is the unique which satisfy to $(x^2, x, x) = 0$ or else, to $(x^2, x^2, x) = 0$.
2. $\mathcal{H}^*$ is the unique which satisfy to $(x, x, x^2) = 0$ or else, to $(x, x^2, x^2) = 0$.
3. $\mathcal{H}^*$ is the unique which satisfy to $(x, x, x) = 0$ or else, to $(x, x^2, x) = 0$. $\square$

We shall have the opportunity to use the supplementary notations $\mathcal{H}_1 = \mathcal{H}$, $\mathcal{H}_2 = {}^*\mathcal{H}$, $\mathcal{H}_3 = \mathcal{H}^*$, $\mathcal{H}_4 = \mathcal{H}^*$. In our computations the first purely imaginary element i in the canonical basis $\{1, i, j, k\}$ of algebra $\mathcal{H}$, will be used.

Now we need the following preliminary result:

Lemma 3.3 *Assume that for $a \in \mathcal{H}$ the equality $[axa, x] = 0$ holds for all $x \in Im(\mathcal{H})$. Then $a \in \mathbb{R}$.*

Proof. For all $x \in Im(\mathcal{H})$, we have

$$\begin{aligned} 0 &= [axa, x] \\ &= [2(\overline{x}|a)a - ||a||^2\overline{x}, x] \quad \text{by triple product identity} \\ &= -2(x|a)[a, x]. \end{aligned}$$

If $a \notin \mathbb{R}$ then the choice of an element $x \in Im(\mathcal{H})$ neither orthogonal nor linearly dependent to $Im(a)$ leads to a contradiction. $\square$

Let us note now [pqrn] the equation deduced from identity $(x^p, x^q, x^r) = 0$, if it occurs, in algebra $\mathcal{H}_n(a, b)$. Theses 32 equations are illustrated in the following table:

Identity $(x^p, x^q, x^r) = 0$	Corresponding equation	[pqrn]
$(x, x, x) = 0$	$ax^2bx = xax^2b$	[1111]
	$\bar{b} \bar{x} \bar{a} x a x = \bar{x} a \bar{x} a x b$	[1112]
	$a x b \bar{x} b \bar{x} = x b x \bar{b} \bar{x} \bar{a}$	[1113]
	$b x \bar{b} x^2 = x^2 \bar{a} x a$	[1114]
$(x, x, x^2) = 0$	$ax^2bax^2 = xaxax^2b$	[1121]
	$\bar{b} \bar{x} \bar{a} x a \bar{x} a x = \bar{x} a \bar{x} a \bar{x} a x b$	[1122]
	$a x b \bar{x} b x = x b a x b \bar{x}$	[1123]
	$\bar{b} x^2 \bar{a} \bar{b} x = \bar{x} \bar{b} a \bar{x}^2 b$	[1124]
$(x, x^2, x) = 0$	$a x a x^2 b^2 x = x a^2 x^2 b x b$	[1211]
	$\bar{b}^2 \bar{x} \bar{a} x \bar{a} x a x = \bar{x} a \bar{b} \bar{x} \bar{a} x a x b$	[1212]
	$a x b x \bar{b} \bar{x} \bar{a} b \bar{x} = x b x \bar{b} x \bar{b} \bar{x} \bar{a}^2$	[1213]
	$\bar{b} a \bar{x}^2 b x \bar{a} \bar{x} = \bar{x} \bar{b} x a \bar{x}^2 b \bar{a}$	[1214]
$(x, x^2, x^2) = 0$	$x a^2 x^2 b a x^2 b = a x a x^2 b^2 a x^2$	[1221]
	$\bar{b}^2 \bar{x} \bar{a} x \bar{a} x a \bar{x} a x = \bar{x} a \bar{b} \bar{x} \bar{a} x a \bar{x} a x b$	[1222]
	$a x b x \bar{b} \bar{x} \bar{a} b x \bar{b} \bar{x} = x b a x b \bar{x} \bar{b} x \bar{b} \bar{x} \bar{a}$	[1223]
	$\bar{b} a \bar{x}^2 b x \bar{a} \bar{b} x^2 = \bar{x} \bar{b} a \bar{x}^2 b a \bar{x}^2 b$	[1224]
$(x^2, x, x) = 0$	$ax^2bxbx = x^2bax^2b$	[2111]
	$x a \bar{x} a x b = \bar{x} a x b a x$	[2112]
	$x b \bar{x} b x \bar{b} \bar{x} \bar{a} = a x b \bar{x} b \bar{x} b \bar{x}$	[2113]
	$x \bar{a} \bar{b} x^2 \bar{a} = a \bar{x}^2 b \bar{a} \bar{x}$	[2114]
$(x^2, x, x^2) = 0$	$ax^2bxbax^2 = x^2baxax^2b$	[2121]
	$\bar{x} a x b a \bar{x} a x = x a \bar{x} a \bar{x} a x b$	[2122]
	$a x b \bar{x} b \bar{x} b x = x b \bar{x} b a x b \bar{x}$	[2123]
	$x \bar{a} \bar{b} a \bar{x}^2 b = a \bar{x}^2 b \bar{a} \bar{b} x$	[2124]
$(x^2, x^2, x) = 0$	$ax^2bax^2b^2x = x^2ba^2x^2bxb$	[2211]
	$\bar{x} \bar{a} x a \bar{b} \bar{x} \bar{a} x a x b = \bar{b} \bar{x} \bar{a} x \bar{a} \bar{x} a x b a x$	[2212]
	$x b \bar{x} b x \bar{b} x \bar{b} \bar{x} \bar{a}^2 = a x b \bar{x} b x \bar{b} \bar{x} \bar{a} b \bar{x}$	[2213]
	$x^2 \bar{a} \bar{b} x a \bar{x}^2 b \bar{a} = a \bar{x}^2 b a \bar{x}^2 b \bar{a} \bar{x}$	[2214]
$(x^2, x^2, x^2) = 0$	$(ax^2b)^2 b a x^2 = x^2 b a (ax^2b)^2$	[2221]
	$\bar{b} \bar{x} \bar{a} x \bar{a} \bar{x} a x b a \bar{x} a x = \bar{x} \bar{a} x a \bar{b} \bar{x} \bar{a} x a \bar{x} a x b$	[2222]
	$a x b \bar{x} b x \bar{b} \bar{x} \bar{a} b x \bar{b} \bar{x} = x b \bar{x} b a x b \bar{x} \bar{b} x \bar{b} \bar{x} \bar{a}$	[2223]
	$(a \bar{x}^2 b)^2 \bar{a} \bar{b} x^2 = x^2 \bar{a} \bar{b} (a \bar{x}^2 b)^2$	[2224]

Remarks 3.4 Let p, q, r be fixed in $\{1, 2\}$ and let $n \in \{1, 2, 3, 4\}$.

1. If $a, b \in \{1, -1\}$, then $\mathcal{H}_n(a, b)$ is isomorphic to $\mathcal{H}_n$.
2. The equation [pqr3] follows from [rqp2] by application of the standard involution and replacement of (a, b) by $(\bar{b}, \bar{a})$.
3. The equation [2q14] follows from [1q24] by application of the standard involution and replacement of (a, b) by $(\bar{b}, \bar{a})$. $\square$

We shall use the following equalities:

$$a = \operatorname{Re}(a) + \operatorname{Im}(a) \quad (3.3)$$

$$\operatorname{Im}(a^3) = (3\operatorname{Re}(a)^2 + \operatorname{Im}(a)^2)\operatorname{Im}(a) \quad (3.4)$$

We have the following preliminary results:

Lemma 3.5 $\mathcal{H}(a, b)$ satisfies to $(x, x^2, x) = 0$ if and only if $a, b \in \{1, -1\}$. In this case $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$.

Proof. The equation [1211] gives $ab = ba$, by putting $x = a$. So $\operatorname{Im}(a)$ and $\operatorname{Im}(b)$ are linearly dependent. Let now x be a non-zero element in $\operatorname{Im}(\mathcal{H})$, orthogonal to a . Then, x is also orthogonal to b , and we have:

$$xa = \bar{a}x, \quad bx = x\bar{b}, \quad xa^2 = \bar{a}^2x, \quad b^2x = x\bar{b}^2.$$

For such an element, the equation [1211] gives $\bar{b}^2 = \bar{a}^2$. As a commutes with b , we have $a = \pm b$ and equation [1211] becomes $[xa^3x, a] = 0$ for all $x \in \operatorname{Im}(\mathcal{H})$. The triple product identity, applied to x, a^3 , gives

$$\operatorname{Re}(xa^3)[x, \operatorname{Im}(a)] = 0 \quad \text{for all } x \in \operatorname{Im}(\mathcal{H}) \quad (3.5)$$

If $\operatorname{Im}(a) \neq 0$, then by putting $x = \operatorname{Im}(a) + v$, where v is a non-zero element in $\operatorname{Im}(\mathcal{H})$, orthogonal to a , the equalities (3.4) and (3.5) give $\operatorname{Im}(a^3) = 0$, that is, $a^3 = \pm 1$. Next, we put $x = 1 + u$ where u is an element in $\operatorname{Im}(\mathcal{H})$, orthonormal to a . We have

$$x^2 = 2u, \quad au = u\bar{a}, \quad a^2u = u\bar{a}^2, \quad aua = u, \quad axa = bxb = a^2 + u.$$

The equation [1211] gives

$$(a^2 + u)ua^2(1 + u) = (1 + u)a^2u(a^2 + u).$$

It follows that $\bar{a}^2 = a^2 \in \mathbb{R}$. Of this fact $a = a^3\bar{a}^2 \in \mathbb{R}$, contradicting the hypothesis $\text{Im}(a) \neq 0$. So $a = \pm b \in \{1, -1\}$ and $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$. $\square$

Lemma 3.6 *Algebra $^*\mathcal{H}(a, b)$ satisfies none of four identities $(x, x, x) = 0$, $(x, x^2, x) = 0$, $(x, x, x^2) = 0$, $(x, x^2, x^2) = 0$.*

Proof. We suppose the opposite.

1. **The first two identities.** We put $x = a$, next $x = \bar{b}$ in [1212], and we get $ab = \bar{b}\bar{a} = \bar{b}a$. It follows that a and b are scalars, and consequently $x^2 = \|x\|^2$ for all $x \in \mathcal{H}$. This absurdity shows that the identity $(x, x^2, x) = 0$, and also $(x, x, x) = 0$, are eliminated.
2. **The third identity.** We put $x = 1$, next $x = a$ in [1122], and we get $\bar{b}a = a^3b = ab$. It follows that $a, b \in \{1, -1\}$. Thus the equality [1122] becomes $\|x\|^4 = \|x\|^2\bar{x}^2$ for all $x \in \mathcal{H}$. Absurd.
3. **The fourth identity.** By putting successively $x = 1$, $x = a$, $x = a^2$ in equation [1222], we get the equalities

$$\bar{b}^2 = a\bar{b}ab \tag{3.6}$$

$$\bar{b}a = ab \tag{3.7}$$

$$\bar{b}^2a^2 = \bar{a}\bar{b}ab \tag{3.8}$$

By taking into account (3.7), the equalities (3.6) and (3.8) give $\bar{b}^2 = a^2b^2$ and $\bar{b}^2a^2 = b^2$, respectively. We deduce that $a^2 = b^4 \in \{1, -1\}$ and we distinguish following both cases:

1. If $a^2 = 1$, the equation [1222] becomes $\|x\|^4\bar{b}^2x = \|x\|^4\bar{x}$ for all $x \in \mathcal{H}$. Absurd.
2. If $a^2 = -1$, then by putting $x = \bar{b}$ in [1222] and taking into account (3.7), we get $b^2 = -1$. Consequently $a^2 = b^4 = 1$. Absurd. $\square$

We get, by means of Remark 3.5 2), a result similar to the Lemma 3.7:

Corollary 3.7 *Algebra $\mathcal{H}^*(a, b)$ satisfies none of four identities $(x, x, x) = 0$, $(x, x^2, x) = 0$, $(x^2, x, x) = 0$, $(x^2, x^2, x) = 0$. $\square$*

We note $a^{\frac{1}{n}}$ a n^{th} root of a in $\mathcal{H}$.

Lemma 3.8 *Algebra $\mathcal{H}^*(a, b)$ satisfies none of four identities $(x, x, x^2) = 0$, $(x, x^2, x^2) = 0$, $(x^2, x, x) = 0$, $(x^2, x^2, x) = 0$.*

Proof. We suppose the opposite.

1. For the first two identities $(x, x^q, x^2) = 0$, we put $x = 1$, next $x = a^{\frac{1}{2}}$ in [1q24], and we get $\bar{a}^2 = b^2 = a$. We deduct from it that a and b commute and $a = \pm \bar{b}$. Thus the equation [1q24] becomes $\bar{b}x^3 = \bar{x}\bar{b}^2\bar{x}^2b$. This gives $b = \pm 1$, by putting $x = \bar{b}^{\frac{1}{3}}$, and leads to $x^3 = \bar{x}^3$. This absurdity shows that both equations [1q24] cannot come true.
2. The Remark 3.5 3) shows that the last two equations [2q14] cannot come true also. $\square$

We have just proved the following result:

Proposition 3.9 *Among those four types of principal isotopes of $\mathcal{H}$*

1. $\mathcal{H}$ is the unique algebra satisfying to all identities of the form $(x^p, x^q, x^r) = 0$.
2. Each of three algebras ${}^*\mathcal{H}(a, b)$, $\mathcal{H}^*(a, b)$, $\mathcal{H}^*(a, b)$, of type different from the first one, can, under certain conditions on the parameters a and b , satisfy both identities $(x^2, x, x^2) = (x^2, x^2, x^2) = 0$. Among these three types of algebras
 - (a) ${}^*\mathcal{H}$ is the unique algebra satisfying to $(x^2, x, x) = 0$, or else $(x^2, x^2, x) = 0$.
 - (b) $\mathcal{H}^*$ is the unique satisfying to $(x, x, x^2) = 0$, or else $(x, x^2, x^2) = 0$.
 - (c) $\mathcal{H}^*$ is the unique satisfying to $(x, x, x) = 0$, or else $(x, x^2, x) = 0$. $\square$

3.2 Specification in case $(x, x^q, x) = 0$

We are going to establish some results which will bring supplementary clarification to the Proposition 3.9. First of all a result completing Lemma 3.6 in the frame of identities of the form $(x, x^q, x) = 0$:

Lemma 3.10 $\mathbb{H}^*(a, b)$ satisfy to $(x, x^2, x) = 0$ if and only if $a, b \in \{1, -1\}$. In this case $\mathbb{H}^*(a, b)$ is isomorphic to $\mathbb{H}^*$.

Proof. By putting $x = a$ in [1214] we get $ab = ba$. Next, we take for x a non-zero element in $Im(\mathbb{H})$, orthogonal to a . The equation [1214] give at first $a^2 = b^2$, that is $b = \pm a$ and consequently $[axa, x] = 0$ for all $x \in Im(\mathbb{H})$. Lemma 3.4 and Remark 3.5 1) conclude. $\square$

We can state the following result:

Theorem 3.11 For a 4-dimensional absolute-valued algebra A , the following are equivalent:

1. A is flexible,
2. A is third power-associative,
3. A satisfies to the identity $(x, x^2, x) = 0$,
4. A is isomorphic to $\mathbb{H}$ or $\mathbb{H}^*$.

Proof. The implications $4 \Rightarrow 1 \Rightarrow 2 \Rightarrow 3$ are obvious. Besides, the implication $3 \Rightarrow 4$ is a consequence of Theorem 2.7, the three Lemmas 3.6, 3.7, 3.11 and Corollary 3.8. $\square$

3.3 Specification in cases $(x^p, x^p, x^r) = 0$ and $(x^p, x^q, x^q) = 0$

We have the following result, for the identity $(x, x, x^2) = 0$:

Theorem 3.12 *For a 4-dimensional absolute-valued algebra A , the following are equivalent:*

1. *A satisfies to the identity $(x, x, x^2) = 0$,*
2. *A is isomorphic to $\mathcal{H}$ or $\mathcal{H}^*$.*

Proof. It is enough to show that the first assertion carry away the second one. Indeed, Lemmas 3.7 and 3.9 show that algebras ${}^*\mathcal{H}(a, b)$ and $\mathcal{H}^*(a, b)$ cannot satisfy to $(x, x, x^2) = 0$. For each one of the two other possibilities $\mathcal{H}_l(a, b)$, where $l \in \{1, 3\}$, the equation [1121] gives $ab = ba$, by putting $x = 1$. So $b \in \text{Lin}\{1, a\}$ and equations [1121], [1123] give $ax = xa$ for all $x \in \text{Im}(\mathcal{H})$, that is, $a = \pm 1$. These equations become $bx^2 = x^2b$ and we have $b = \pm 1$. In this case $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$, and $\mathcal{H}^*(a, b)$ is isomorphic to $\mathcal{H}^*$, by virtue of the Remark 3.5 1). $\square$

We deduct from the Theorem 3.13 and Remark 3.5 2) the following result for identity $(x^2, x, x) = 0$:

Theorem 3.13 *For a 4-dimensional absolute-valued algebra A , the following are equivalent:*

1. *A satisfies to $(x^2, x, x) = 0$,*
2. *A is isomorphic to $\mathcal{H}$ or ${}^*\mathcal{H}$. $\square$*

We have the following result, for identity $(x, x^2, x^2) = 0$:

Theorem 3.14 *For a 4-dimensional absolute-valued algebra A , the following are equivalent:*

1. *A satisfies to $(x, x^2, x^2) = 0$,*
2. *A is isomorphic to $\mathcal{H}$, $\mathcal{H}^*$ or $\mathcal{H}^*(1, i)$.*

Proof. We check easily that algebra $\mathcal{H}^*(1, i)$ satisfies to $(x, x^2, x^2) = 0$ and it is enough to show that the first assertion carry away the second one. Indeed, both Lemmas **3.7** and **3.9** show that the algebras ${}^*\mathcal{H}(a, b)$ and $\mathcal{H}^*(a, b)$ cannot satisfy to $(x, x^2, x^2) = 0$. For each of two other possibilities $\mathcal{H}_l(a, b)$, where $l \in \{1, 3\}$, the equation [1221] gives $ab = ba$, by putting $x = 1$. So $b \in \text{Lin}\{1, a\} := E$ and we note $E^\perp$ the subspace of $\mathcal{H}$ orthogonal to E . We distinguish following both cases:

1. If $A = \mathcal{H}(a, b)$. The same arguments as those in the proof of Theorem **3.13** show that $a, b \in \{1, -1\}$. So $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$.
2. If $A = \mathcal{H}^*(a, b)$. The equation [1223] gives $b^4 = a^2$, by putting for x a non-zero element in $E^\perp$, so $a = \pm b^2$. The equation [1223] becomes

$$b^2 x b x \bar{b} \bar{x} \bar{b} x \bar{b} \bar{x} = x b^3 x b \bar{x} \bar{b} x \bar{b} \bar{x} b^2 \quad \text{for all } x \in \mathcal{H} \quad (3.9)$$

By putting $y = bx$, the equality (3.9) becomes

$$b y^2 \bar{b} \bar{y} \bar{b} y \bar{b} \bar{y} b = \bar{b} y b^2 y b \bar{y} \bar{b} y \bar{b} \bar{y} b \quad \text{for all } y \in \mathcal{H} \quad (3.10)$$

We multiply to the right and to the left, by b , the members of the equality (3.10) and we take, for y , a non-zero element in $\text{Im}(\mathcal{H})$. We get, after a simplification to the left by y , the equality $y b y \bar{b} y \bar{b} y b^2 = b^2 y b y \bar{b} y \bar{b} y$, that is

$$[y b y \bar{b} y \bar{b} y, b^2] = 0 \quad \text{for all } y \in \text{Im}(\mathcal{H}) \quad (3.11)$$

We compute now, by means of the triple product identity, the expression $y b y \bar{b} y \bar{b} y$:

$$\begin{aligned} y b y \bar{b} y \bar{b} y &= y b y (2(\bar{y} | \bar{b}) \bar{b} - \bar{y}) y \\ &= y b (2(\bar{y} | \bar{b}) y \bar{b} y - \|y\|^2 y) \\ &= y b \left((4(y | b)^2 - \|y\|^2) y - 2\|y\|^2 (y | b) b \right) \\ &= (4(y | b)^2 - \|y\|^2) y b y - 2\|y\|^2 (y | b) y b^2 \\ &= (4(y | b)^2 - \|y\|^2) (2(\bar{b} | y) y - \|y\|^2 \bar{b}) - 2\|y\|^2 (y | b) y b^2 \\ &= (\|y\|^2 - 4(y | b)^2) (2(b | y) y + \|y\|^2 \bar{b}) - 2\|y\|^2 (b | y) y b^2 \end{aligned}$$

Now, for $y = \text{Im}(b) + u$ where $u \in E^\perp - \{0\}$, we have

$$\begin{aligned} (y | b) &= (\text{Im}(b) + u | b) \\ &= (\text{Im}(b) | b) \\ &= \|\text{Im}(b)\|^2 \end{aligned}$$

and

$$||y||^2 - 4(y|b)^2 = ||Im(b)||^2 + ||u||^2 - 4||Im(b)||^4.$$

In the other hand, we can choose u rather big in such a way that

$$||y||^2 - 4(y|b)^2 := \lambda \neq 0.$$

Thus

$$\begin{aligned} [yby\bar{b}y\bar{b}y, b^2] &= [2||Im(b)||^2(\lambda y - ||y||^2 y b^2) + \lambda ||y||^2 \bar{b}, b^2] \\ &= 2||Im(b)||^2[\lambda y - ||y||^2 y b^2, b^2] \\ &= 2||Im(b)||^2[\lambda u - ||y||^2 u b^2, b^2] \\ &= 2||Im(b)||^2(\lambda(u b^2 - b^2 u) - ||y||^2(u b^4 - b^2 u b^2)) \\ &= 2||Im(b)||^2(\lambda u(b^2 - \bar{b}^2) - ||y||^2 u(b^4 - 1)) \\ &= 2||Im(b)||^2 u(\lambda(b^2 - \bar{b}^2) - ||y||^2(b^4 - 1)) \\ &= 2||Im(b)||^2 u(\lambda \bar{b}^2 - ||y||^2)(b^4 - 1). \end{aligned}$$

Now

$$\begin{aligned} [yby\bar{b}y\bar{b}y, b^2] = 0 &\Leftrightarrow ||Im(b)||^2(\lambda \bar{b}^2 - ||y||^2)(b^4 - 1) = 0 \\ &\Leftrightarrow Im(b) = 0, b^2 = \lambda ||y||^{-2} \text{ or } b^4 = 1 \\ &\Leftrightarrow b^4 = 1. \end{aligned}$$

If $b = \pm 1$, then $A = \mathcal{H}^*(\pm 1, \pm 1)$ is isomorphic to $\mathcal{H}^*$. If $b \in S(Im(\mathcal{H}))$, then b is conjugated to i by Lemma 2.10 and so $A = \mathcal{H}^*(\pm 1, b)$ is isomorphic to $\mathcal{H}^*(1, i)$ by virtue of Theorem 2.7. $\square$

We deduce from the Theorem 3.14 and Remark 3.5 2) the following result for the identity $(x^2, x^2, x) = 0$:

Theorem 3.15 *For a 4-dimensional absolute-valued algebra A , the following are equivalent:*

1. *A satisfy to $(x^2, x^2, x) = 0$,*
2. *A is isomorphic to $\mathcal{H}$, ${}^*\mathcal{H}$ or ${}^*\mathcal{H}(i, 1)$.* $\square$

3.4 Specification in case $(x^2, x^q, x^2) = 0$

We are now going to determine those 4-dimensional absolute-valued algebras which satisfy to an identity of the form $(x^2, x^q, x^2) = 0$. We shall see that there is a variety, among the algebras $\mathcal{H}_2(a, b)$ and $\mathcal{H}_3(a, b)$, satisfying such an identity. Let us begin, at first, with the algebras $\mathcal{H}_1(a, b)$ and $\mathcal{H}_4(a, b)$:

Lemma 3.16 *The following are equivalent:*

1. $\mathcal{H}(a, b)$ satisfy to $(x^2, x, x^2) = 0$.
2. $\mathcal{H}(a, b)$ satisfy to $(x^2, x^2, x^2) = 0$.
3. $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$.

Proof. It is enough to show that the third assertion is consequence of each of both first ones.

1. **1 $\Rightarrow$ 3.** The equation [2121] gives $abxba = baxab$ for all $x \in \mathcal{R} \cup \text{Im}(\mathcal{H})$. So

$$abxba = baxab \quad \text{for all } x \in \mathcal{H} \quad (3.12)$$

We deduce from it that $ba\bar{b}\bar{a} = \bar{b}\bar{a}ba$ is a central element of $\mathcal{H}$, and we have $ab = \pm ba$. Besides, the case $ab = -ba$ is eliminated at once by making $x = b^{\frac{1}{2}}$ in [2121]. So $ab = ba$ et b belongs in the subalgebra $\mathcal{R}_H[a] = \text{Lin}\{1, a\}$. Let now u be an element in $\text{Im}(\mathcal{H})$ orthonormal to a , then u is orthogonal to $\text{Lin}\{1, a\}$. By putting $x = 1 + u$ in [2121] and taking into account that the $\{a, b, \bar{a}, \bar{b}\}$ is a commutative set and $au = u\bar{a}, ub = \bar{b}u$, we get $\bar{a}(1 + u)b = a(1 + u)\bar{b}$ or still

$$(1 + u)b^2 = a^2(1 + u) \quad (3.13)$$

As $a^2u = u\bar{a}^2$, we have $b^2 - a^2 = u(\bar{a}^2 - b^2)$. The trace property of $(\cdot | \cdot)$, and the fact that $b^2 - a^2, \bar{a}^2 - b^2 \in \text{Lin}\{1, a\}$, show that $b^2 - a^2$ and $u(\bar{a}^2 - b^2)$ are orthogonal to each other. So $b^2 = a^2 = \bar{a}^2$, and, taking into account that a and b commute, we have $b = \pm \bar{a}$. The Remarks 2.8 1), 2.8 3) and ([CR 08] Corollary 3.2) show then that $\mathcal{H}(a, b)$ is isomorphic to $\mathcal{H}$.

2. **2 $\Rightarrow$ 3.** The equation [2221] gives $ab = ba$, by taking, for x , a square root of $\bar{a}$. So $\text{Im}(a), \text{Im}(b)$ are linearly dependent. By putting $x = 1 + u$ for norm-one u in $\text{Im}(\mathcal{H})$, orthogonal to a , we get $x^2 = 2u, ua = \bar{a}u$ and $ub = \bar{b}u$. For such an element x , the equation [2221] gives $ab = \bar{a}\bar{b}$. As $ab = ba$, we have $b = \pm \bar{a}$. $\square$

Lemma 3.17 *The following are equivalent:*

1. $\overset{*}{\mathbb{H}}(a, b)$ satisfy to $(x^2, x, x^2) = 0$.
2. $\overset{*}{\mathbb{H}}(a, b)$ satisfy to $(x^2, x^2, x^2) = 0$.
3. $\overset{*}{\mathbb{H}}(a, b)$ is isomorphic to $\overset{*}{\mathbb{H}}$.

Proof. It is enough to show that the third assertion is consequence of each of both first ones.

1. **1** $\Rightarrow$ **3.** The equation [2124] gives $x\bar{a}bab = ab\bar{a}bx$ for all $x \in \mathbb{R} \cup Im(\mathbb{H})$. So

$$x\bar{a}bab = ab\bar{a}bx \quad \text{for all } x \in \mathbb{H} \quad (3.14)$$

It follows easily from (3.14) that $\bar{a}bab = ab\bar{a}b \in \mathbb{R}$ and then $ab = \pm ba$. A calculation similar to that of the first part in the proof of Lemma 3.17 ends by $b = \pm \bar{a}$.

2. **2** $\Rightarrow$ **3.** The same as that of the second part in the proof of Lemma 3.17. $\square$

We need, for the identity $(x^2, x, x^2) = 0$, the following supplementary preliminary results:

Lemma 3.18 $\overset{*}{\mathbb{H}}(a, b)$ satisfy to $(x^2, x, x^2) = 0$ if and only if $a = \pm 1$ and $b \in S(\mathbb{H})$ is arbitrary.

Proof. The condition $a = \pm 1$ is sufficient, according to the equation [2122], and it is enough to show that it is necessary. Indeed, the equation [2122] gives

$$[b, (ax)^2] = 0 \quad \text{for all } x \in Im(\mathbb{H}) \quad (3.15)$$

Next, we put $x = Im(a) + \bar{a}u$, where u is an element in $Im(\mathbb{H})$ orthogonal to a , and we have

$$(ax)^2 = Im(a)^4 + Re(a)^2 Im(a)^2 + u^2 + 2Im(a)^2 (Re(a)Im(a) + u) \quad (3.16)$$

If $Im(a) \neq 0$, then the equalities (3.15) and (3.16) given $[b, Re(a)Im(a) + u] = 0$ for all $u \in Im(\mathbb{H})$ orthogonal to a . Consequently b is a scalar, and the equation [2122] give $[Im(a), x^2] = 0$ for all $x \in \mathbb{H}$. Absurd. $\square$

As a consequence of Lemma 3.18 and Remark 3.5 2):

Lemma 3.19 $\mathcal{H}^*(a, b)$ satisfy to $(x^2, x, x^2) = 0$ if and only if $b = \pm 1$ and $a \in S(\mathcal{H})$ is arbitrary. $\square$

We have just proved, by means of Lemma 2.10, the following result:

Theorem 3.20 For a 4-dimensional absolute-valued algebra A , the following are equivalent:

1. A satisfy to identity $(x^2, x, x^2) = 0$,
2. A is isomorphic to $\mathcal{H}$, $\mathcal{H}^*$, ${}^*\mathcal{H}(1, e^{i\alpha})$ or $\mathcal{H}^*(e^{i\alpha}, 1)$ with $\alpha \in [0, \pi[$. $\square$

We need, for the identity $(x^2, x^2, x^2) = 0$, the following supplementary preliminary result:

Lemma 3.21 ${}^*\mathcal{H}(a, b)$ satisfies to $(x^2, x^2, x^2) = 0$ if and only if one of the following conditions holds

1. $a^2 = 1$ and $b \in S(\mathcal{H})$. In this case ${}^*\mathcal{H}(a, b)$ is isomorphic to ${}^*\mathcal{H}(1, e^{i\alpha})$ where $\alpha \in [0, \pi[$.
2. $a^2 = -1$ and $b \in \{1, -1, a, -a\}$. In this case ${}^*\mathcal{H}(a, b)$ is isomorphic to ${}^*\mathcal{H}(i, 1)$ or ${}^*\mathcal{H}(i, i)$.

Proof. The equation [2222] is obviously verified if one of both conditions 1 or 2 holds. Let us suppose now that ${}^*\mathcal{H}(a, b)$ satisfy to $(x^2, x^2, x^2) = 0$.

1. We show that $a^4 = 1$. Indeed, the equation [2222] gives at once

$$a^2b = ba^2 \tag{3.17}$$

by putting $x = a$. Now, for norm-one x in $Im(\mathcal{H})$, orthogonal to a and b , we have $xa = \bar{a}x$ and $x\bar{a} = ax$. The equation [2222] gives $\bar{a}^2b = ba^2$ and the condition $a^4 = 1$ is obtained from equality (3.17).

2. If $a^2 = 1$, then ${}^*\mathcal{H}(a, b)$ is isomorphic to ${}^*\mathcal{H}(1, b)$ by virtue of Remark 2.8 1).
3. If $a^2 = -1$, then the equation [2222] is equivalent to

$$\bar{b}\bar{x}(axa)\bar{x}(ax.b\bar{a}x.ax) \in Im(\mathcal{H}) \quad \text{for all } x \in \mathcal{H}. \tag{3.18}$$

- (a) Let us show that $ba = \pm ab$. Indeed, by applying the triple product identity to the expression $ax(ba\bar{x})ax$, for an element x in $\mathcal{H}$ orthogonal to a , we have

$$\begin{aligned}
\bar{x}(axa)\bar{x}(ax.ba\bar{x}.ax) &= \bar{x}^3(2(xa\bar{b}|ax)ax - ||x||^2xa\bar{b}) \\
&= \bar{x}^3(2(a\bar{x}\bar{b}|ax)ax - ||x||^2a\bar{x}\bar{b}) \\
&= \bar{x}^3a(2(\bar{x}\bar{b}|x)x - ||x||^2\bar{x}\bar{b}) \\
&= ax^3(x(bx)x) \quad \text{by triple product identity} \\
&= ax^4bx^2.
\end{aligned}$$

We note E the subspace of $\mathcal{H}$ orthogonal to $\text{Lin}\{1, a\}$. By putting $x = 1 + u$ where $u \in S(E)$, we get $x^2 = 2u$, $x^4 = -4$. The equation [2222] gives $\bar{b}ax^4bx^2 \in \text{Im}(\mathcal{H})$, or else, $\bar{b}abu \in \text{Im}(\mathcal{H})$ for all $u \in E$. The scalar part $(\bar{b}ab|u)$ of $\bar{b}abu$ vanish for all $u \in E$, that is, $\bar{b}ab \in \text{Lin}\{1, a\}$. There are then scalars ν, ρ for which $\bar{b}ab = \nu + \rho a$, or else, $ab = \nu b + \rho ba$. We have

$$\begin{aligned}
0 &= (a|1)(b|b) \\
&= (ab|b) \\
&= (\nu b + \rho ba|b) \\
&= \nu.
\end{aligned}$$

So $ab = \pm ba$.

- (b) Let us show that $b \in \{1, -1, a, -a\}$. Indeed, for any norm-one element x in E , we put

$$x = 1 + \frac{a + u}{\sqrt{2}}$$

and we get:

$$\bar{x}ax = a + u + \sqrt{2}au \quad (3.19)$$

and

$$\bar{x}ax.a.\bar{x}ax = 2a - 2u - 2\sqrt{2}au \quad (3.20)$$

Now, by putting

$$(y, z) = (\bar{b}(a - u - \sqrt{2}au)b, -1 - au + \sqrt{2}u),$$

and taking into account to the equalities (3.19), (3.20) the equation [2222] gives

$$yz = \bar{z}y \quad (3.21)$$

If $ba = -ab$, then $b \in E$, and we choose u to be orthogonal to b . A first computation gives $y = -a + u - \sqrt{2}au$ and the equality (3.21) leads to the absurdity $2\sqrt{2} = -2\sqrt{2}$. So $ab = ba$ and the equality (3.21) gives $\bar{b}^2 = b^2$, or else, $b \in \{1, -1, a, -a\}$. Remark 2.8 1), Lemma 2.10 and Theorem 2.7 conclude. $\square$

As consequence of Lemmas 2.10, 3.21 and Remarks 2.8 1), 3.5 2):

Lemma 3.22 $\mathcal{H}^*(a, b)$ satisfy to $(x^2, x^2, x^2) = 0$ if and only if one the following conditions holds

1. $b^2 = 1$ and $a \in S(\mathcal{H})$. In this case $\mathcal{H}^*(a, b)$ is isomorphic to $\mathcal{H}^*(e^{i\alpha}, 1)$ where $\alpha \in [0, \pi[$.
2. $b^2 = -1$ and $a \in \{1, -1, b, -b\}$. In this case $\mathcal{H}^*(a, b)$ is isomorphic to $\mathcal{H}^*(1, i)$, or $\mathcal{H}^*(i, i)$. $\square$

We have the following result:

Theorem 3.23 For a 4-dimensional absolute-valued algebra A , the following are equivalent:

1. A satisfy to $(x^2, x^2, x^2) = 0$,
2. A is isomorphic to

$$\mathcal{H},$$

$${}^*\mathcal{H}(i, 1), {}^*\mathcal{H}(i, i), {}^*\mathcal{H}(1, e^{i\alpha}), \mathcal{H}^*(1, i), \mathcal{H}^*(i, i), \mathcal{H}^*(e^{i\alpha}, 1) \text{ where } \alpha \in [0, \pi[$$

$$\text{or } {}^*\mathcal{H}.$$

Proof. Consequence of Theorem 2.7 and Lemmas 3.16, 3.17, 3.21, 3.22. $\square$

We summarize the results established in this section as follows:

Theorem 3.24 *Let A be a 4-dimensional AAV. Then A satisfy to an identity of the form $(x^p, x^q, x^r) = 0$, for fixed integers $p, q, r \in \{1, 2\}$, if and only if A is isomorphic to one of principal isotopes of $\mathbb{H}$ described according to the following table:*

<i>A satisfy to</i>	<i>The list of isomorphism classes</i>
$(x, x, x) = 0$ or $(x, x^2, x) = 0$	$\mathbb{H}, \mathbb{H}^*$
$(x, x, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*$
$(x, x^2, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*, \mathbb{H}^*(1, i)$
$(x^2, x, x) = 0$	$\mathbb{H}, {}^*\mathbb{H}$
$(x^2, x^2, x) = 0$	$\mathbb{H}, {}^*\mathbb{H}, {}^*\mathbb{H}(i, 1)$
$(x^2, x, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*$ ${}^*\mathbb{H}(1, e^{i\alpha}), \mathbb{H}^*(e^{i\alpha}, 1)$ with $\alpha \in [0, \pi[$
$(x^2, x^2, x^2) = 0$	$\mathbb{H}, \mathbb{H}^*$ ${}^*\mathbb{H}(i, 1), {}^*\mathbb{H}(i, i), \mathbb{H}^*(1, i), \mathbb{H}^*(i, i)$ ${}^*\mathbb{H}(1, e^{i\alpha}), \mathbb{H}^*(e^{i\alpha}, 1)$ with $\alpha \in [0, \pi[$

Remarks 3.25 .

1. All isomorphism classes, in above table, are represented by commuting parameters a, b . Theorem 2.9 shows that every 4-dimensional absolute-valued algebra satisfying to an identity of the form $(x^p, x^q, x^r) = 0$, for fixed p, q, r in $\{1, 2\}$, contains 2-dimensional subalgebras. Concretely, algebras $\mathbb{H}$ and $\mathbb{H}^*$ contain $\mathbb{C}$ and $\mathbb{C}^*$, respectively. Those algebras ${}^*\mathbb{H}(a, b)$ and $\mathbb{H}^*(a, b)$, in above table, contain ${}^*\mathbb{C}$ and $\mathbb{C}^*$, respectively.
2. Among those 4-dimensional absolute-valued algebras, each identity of the form $(x^p, x^q, x^r) = 0$, other than $(x^2, x^2, x^2) = 0$, carry away $(x^2, x^2, x^2) = 0$. The converse is false.
3. Every 4-dimensional absolute-valued algebra satisfying any one of identities $(x, x, x) = 0$, $(x, x^2, x) = 0$, $(x, x, x^2) = 0$ or $(x^2, x, x) = 0$ cannot be single-generated. $\square$

4 Variety in dimension eight for absolute-valued algebras satisfying $(x^2, y, x^2) = 0$

4.1 Possibility of absence of 4-dimensional subalgebras

In this section we are going to put evidence the unlimitedness of the class of 8-dimensional absolute-valued algebras satisfying to $(x^2, y, x^2) = 0$ by giving examples of such an algebras which contain no 4-dimensional subalgebras.

Gleichgewicht [G 63] defined on the normed space, underlying to an absolute-valued algebra $(A, ||.||)$ with involution $*$, a new product $\odot$ by putting $x \odot y = x^*y$ for all $x, y \in A$. This provides a new absolute-valued algebra $(A, \odot, ||.||) := A_*$ called a cracovian algebra generated by A . Besides, there exists an element $e \in A$, such that $x^*x = ||x||^2e$ for all $x \in A$ [G 63]. This shows that e is the unique non-zero idempotent for cracovian algebra A_* .

Now, let us remind that every absolute-valued algebra $(A, ||.||)$ containing a non-zero central idempotent is endowed with an involution [Elm 90] and that the norm $||.||$ comes from an inner product $(.|.)$ [U 61]. It is shown that if $(A, ||.||, (.|.))$ is an absolute-valued real algebra which is not isomorphic to $\mathcal{C}^*$, containing a non-zero central e , admits one and only one involution $*$ ([RR 09] Corollary 3.8), given by $x \mapsto x^* := 2(x|e)e - x$ ([RR 09] Proposition 3.5). It follows immediately the result:

Proposition 4.1 *Let A be an absolute valued real algebra which is not isomorphic to $\mathcal{C}^*$, containing a non-zero central idempotent e , and let $*$ its unique involution. Then every subspace of A containing e is $*$ -invariant. $\square$*

For every linear isometry f of euclidian space $\mathcal{O}$, fixing 1, the normed space $\mathcal{O}$ endowed with the product $\bullet$ defined, for all $x, y \in \mathcal{O}$, by $x \bullet y = f(x)f(y)$ becomes an absolute-valued real algebra $\mathcal{O}^f$ with central idempotent 1 [RR 09]. According to ([RR 09] Proposition 3.5) the mapping $\alpha.1 + u \mapsto^* \alpha.1 - u$ is an involution on algebra $\mathcal{O}^f = \mathbb{R}.1 \oplus Im(\mathcal{O})$. In the other hand, it is well known that any finite-dimensional absolute-valued algebra contains a non-zero idempotent [Se 54]. With the same notations as above, we have the following result:

Proposition 4.2 *Every subalgebra B of $(\mathcal{O}^f)_*$ is a subalgebra of $\mathcal{O}^f$.*

Proof. The unique non-zero idempotent $e = 1$ of algebra $(\mathcal{O}^f)_*$ belongs to B . So the subspace B of $\mathcal{O}$ is $*$ -invariant by virtue of Proposition 4.1, and then B is a subalgebra of $\mathcal{O}^f$. $\square$

We have the following result about a flexible idempotent:

Proposition 4.3 *Let A be an absolute-valued algebra A of finite-dimension $n \geq 2$, having a nonzero flexible idempotent e . Then A contains a 2-dimensional subalgebra invariant under L_e and R_e .*

Proof. The operators L_e, R_e are linear isometries fixing e , and induce isometries on the orthogonal space $(\mathbb{R}e)^\perp := E$. As E has odd dimension and L_e commutes with R_e , there exists common norm-one eigenvector $u \in E$ for both L_e and R_e associated to eigenvalues $\alpha, \beta \in \{1, -1\}$. Now, the normed space A endowed with multiplication $x \odot y = L_e(x)R_e(y)$ is an absolute-valued algebra for which we check easily that e is a central idempotent. So $u \odot u = -e$ ([Elm 90] Lemma 3.3), that is, $u^2 = -\alpha\beta e$. It follows that $\text{Lin}\{e, u\}$ is a 2-dimensional subalgebra of A invariant under both L_e and R_e . $\square$

Remark 4.4 *There exists an appropriate linear isometry f , of euclidian space $\mathcal{O}$, for which the algebra $\mathcal{O}^f$ contains no 4-dimensional subalgebras ([RR 09] Theorem 4.8, Corollary 4.9). $\square$*

A finite-dimensional algebra A over a field K is said to be of degree $n \in \mathbb{N}$ if n is the minimum natural number such that all single-generated subalgebras of A have dimension $\leq n$. It follows from Theorem 2.5, that finite-dimensional absolute valued real algebras have degree 1, 2, 4 or 8. Rodríguez ([Rod 94] Theorem 3) showed that $\mathcal{C}, {}^*\mathcal{C}, \mathcal{C}^*, {}^*\mathcal{C}^*, \mathcal{H}, {}^*\mathcal{H}, \mathcal{H}^*, {}^*\mathcal{H}^*, \mathcal{O}, {}^*\mathcal{O}, \mathcal{O}^*, {}^*\mathcal{O}^*$ (the standard isotopes of $\mathcal{C}, \mathcal{H}, \mathcal{O}$) and $\mathcal{P}$ (the pseudo-octonion algebra [Ok 78]) are the only absolute valued real algebras of degree 2.

Remarks 4.5 .

1. *Every finite-dimensional third power-associative absolute valued real algebra has degree ≤ 2 ([Elm 87], [EP 94]).*
2. *Every 8-dimensional absolute valued real algebra of degree 2 contains 4-dimensional subalgebras.* $\square$

Now, with the same notation as in Remark 4.4, we state the following main result of this subsection:

Proposition 4.6 *Let B denotes algebra $(\mathcal{O}^f)_*$. Then*

1. *B contains a 2-dimensional subalgebra but no 4-dimensional subalgebras. As immediate consequence, B has degree 8.*
2. *B satisfy to $(x^2, y, x^2) = 0$ but do not satisfy to $(x, x^2, x) = 0$.*

Proof.

1. A direct computation shows that $e = 1 \in B$ is a flexible idempotent, so B contains a 2-dimensional subalgebra according to Proposition 4.3. In other hand B contains no 4-dimensional subalgebras by virtue of Remark 4.4 and Proposition 4.2. By Remark 4.5.2 B has degree 8.
2. Follows by direct computation. $\square$

4.2 On 8-dimensional absolute-valued algebras satisfying $(x, x^2, x) = 0$

Among those 8-dimensional absolute-valued algebras satisfying $(x, x^2, x) = 0$ we give sufficient conditions which insure third power-associativity:

Theorem 4.7 *Let A be a finite-dimensional absolute-valued algebra satisfying $(x, x^2, x) = 0$, then the degree of A is $\neq 4$. Moreover, the following are equivalent:*

1. A is flexible,
2. A is third power-associative,
3. A has degree ≤ 2 ,
4. A has degree $\neq 8$,
5. A is isomorphic to $\mathbb{R}, \mathbb{C}, \mathbb{C}^*, \mathbb{H}, \mathbb{H}^*, \mathbb{O}, \mathbb{O}^*$, or $\mathbb{P}$.

Proof. The first proposition is a consequence of Remark 3.25.3. This shows the equivalence $\mathbf{3} \Leftrightarrow \mathbf{4}$. In the other hand, the fifth assertion carry away all four above ones. Moreover $\mathbf{1} \Rightarrow \mathbf{2}$ is clear. Now, the implication $\mathbf{2} \Rightarrow \mathbf{3}$ follows from Remark 4.5.1. Finally $\mathbf{3} \Rightarrow \mathbf{5}$ follows by computation using [Rod 94]. This completes the proof. $\square$

Problem 4.8 *Let A be an 8-dimensional absolute-valued real algebra satisfying both identities $(x, x^2, x) = (x^2, y, x^2) = 0$. It is interesting to know if algebra A is third power associative, or even contains 4-dimensional subalgebras.* $\square$

References

- [1] [A 47] A. A. Albert, *Absolute valued real algebras*. Ann. Math. **48**, (1947) 495-501.
- [2] [BBO 82] G. M. Benkart, D. J. Britten, and J. M. Osborn, *Real flexible division algebras*. Can. J. Math. XXXIV **3**, (1982) 550-588.
- [3] [CKMMRR] A. Calderón, A. Kaidi, C. Martín, A. Morales, M. Ramírez, and A. Rochdi, *Finite-dimensional absolute valued algebras*. (To appear in "Israel Journal of Mathematics")
- [4] [CM 05] A. Calderón and C. Martín, *Two-graded absolute-valued algebras*. J. Algebra **292** (2005), 492-515.

- [5] [CR 08] A. Chandid and A. Rochdi, *A survey on absolute valued algebras satisfying $(x^i, x^j, x^k) = 0$* . Int. J. Algebra, **2**, (2008) 837-852.
- [6] [Cu 06] J. A. Cuenca, *On composition and absolute valued algebras*. Proceedings of the Royal Society of Edinburgh, **136** A, (2006) 717-731.
- [7] [EP 94] A. Elduque and J. M. Pérez, *Third power associative composition algebras*. Manuscripta Math. **84**, (1994) 73-87.
- [8] [EP 99] A. Elduque and J. M. Pérez, *Composition algebras of degree two*. Proc. Edinburgh. Math. Soc. **42** (1999), 641-653.
- [9] [Elm 82] M. L. El-Mallah, *Quelques résultats sur les algèbres absolument valuées*. Arch. Math., **38** (1982), 432-437.
- [10] [Elm 83] M. L. El-Mallah, *Sur les algèbres absolument valuées qui vérifient l'identité $(x, x, x) = 0$* . J. Algebra **80** (1983), 314-322.
- [11] [Elm 87] M. L. El-Mallah, *On finite dimensional absolute valued algebras satisfying $(x, x, x) = 0$* . Arch. Math. **49** (1987), 16-22.
- [12] [Elm 88] M. L. El-Mallah, *Absolute valued algebras with an involution*. Arch. Math. **51**, (1988), 39-49.
- [13] [Elm 90] M. L. El-Mallah, *Absolute valued algebras containing a central idempotent*. J. Algebra **128** 180-187 (1990).
- [14] [Elm 97] M. L. El-Mallah, *Absolute valued algebraic algebras satisfying $(x, x, x) = 0$* . Pure Math. Appl. **8** (1997), 39-52.
- [15] [Elm 01] M. L. El-Mallah, *Absolute valued algebras satisfying $(x, x, x^2) = 0$* . Arch. Math. **77** (2001), 378-382.
- [16] [EE 04] M. L. El-Mallah and M. El-Agawany, *Absolute valued algebras satisfying $(x^2, x^2, x^2) = 0$* . Comm. Algebra **32**, 3537-3541 (2004).
- [17] [EM 80] M. L. El-Mallah and A. Micali, *Sur les algèbres normées sans diviseurs topologiques de zéro*. Boletín de la Sociedad Matemática Mexicana **25**, (1980), 23-28.
- [18] [EM 81] M. L. El-Mallah and A. Micali, *Sur les dimensions des algèbres absolument valuées*. J. Algebra **68**, (1981), 237-246.

- [19] [HKR 91] F. Hirzebruch, M. Koecher and R. Remmert, *Numbers*. Springer-Verlag, (1991).
- [20] [M 38] S. Mazur, *Sur les anneaux linéaires*. CRAS Paris **207** (1938), 1025-1027.
- [21] [MRR] A. Morales, M. I. Ramírez, and A. Rochdi, *Eight-dimensional absolute-valued algebras with left-unit satisfying $(x^2, x^2, x^2) = 0$* . (Preprint).
- [22] [Ok 78] S. Okubo, *Pseudo-quaternion and pseudo octonion algebras*. Hadronic J. **1**, (1978) 1250-1278.
- [23] [Osb 62] J. M. Osborn, *Quadratic division algebras*. Trans. AMS **105** (1962), 202-221.
- [24] [Ost 18] A. Ostrowski, *Über einige Lösungen der Funktionalgleichung $\Psi(x)\Psi(x) = \Psi(xy)$* . Acta Math. **41**, (1918) 271-284.
- [25] [Ra 99] M. I. Ramírez, *On four dimensional absolute valued algebras*. Proceedings of the International Conference on Jordan Structures (Málaga, June 1997) (Ed. A. Castellón, J. A. Cuenca, A. Fernández, and C. Martín), 169-173, Málaga, (1999)
- [26] [Roc 03] A. Rochdi, *Eight-dimensional real absolute valued algebras with left unit whose automorphism group is trivial*. IJMMS **70**, (2003) 4447-4454.
- [27] [RR 09] A. Rochdi and A. Rodríguez, *Absolute valued algebras with involution*. Communications in Algebra **37**, (2009) 1151-1159.
- [28] [Rod 94] A. Rodríguez, *Absolute valued algebras of degree two*. In Nonassociative Algebra and its applications (Ed. S. González), Kluwer Academic Publishers, Dordrecht-Boston-London (1994), 350-356.
- [29] [Sc 66] R. D. Schafer, *An introduction to nonassociative algebras*. Academic Press, New York (1966).
- [30] [Se 54] B. Segre, *La teoria delle algebre ed alcune questione di realta*. Univ. Roma, Ist. Naz. Alta. Mat., Rend. Mat. E Appl. Serie 5, **13** (1954), 157-188.
- [31] [UW 60] K. Urbanik and F. B. Wright, *Absolute valued algebras*. Proc. Amer. Math. Soc. **11** (1960), 861-866.

DÉPARTEMENT DE MATHÉMATIQUES ET INFORMATIQUE,
FACULTÉ DES SCIENCES BEN M'SIK, UNIVERSITÉ HASSAN II-MOHAMMEDIA,
B.P. 7955 CASABLANCA (MOROCCO)

E-mail address: ahchandid@hotmail.com

DEPARTAMENTO DE ÁLGEBRA Y ANÁLISIS MATEMÁTICO,
UNIVERSIDAD DE ALMERÍA. 04071 ALMERÍA (SPAIN)

E-mail address: mramirez@ual.es

DÉPARTEMENT DE MATHÉMATIQUES ET INFORMATIQUE,
FACULTÉ DES SCIENCES BEN M'SIK, UNIVERSITÉ HASSAN II-MOHAMMEDIA,
B.P. 7955 CASABLANCA (MOROCCO)

E-mail address: abdellatifro@yahoo.fr